

This question paper contains 4 printed pages]

Roll No.

--	--	--	--	--	--	--	--	--	--	--

S. No. of Question Paper : 2443

Unique Paper Code : 2352401

F-4

Name of the Paper : Linear Algebra

Name of the Course : B.A./B.Sc. (H) — Allied Course

[For the Students of other than Mathematics]

Semester : IV

Duration : 3 Hours

Maximum Marks : 75

*(Write your Roll No. on the top immediately on receipt of this question paper.)*

Attempt *All* questions by selecting any *two* parts from each question.

1. (a) A woman rowing on a wide river wants the resultant (net) velocity of her boat to be 8 km/hr westward. If the current is moving 2 km/hr northeastward, what velocity vector should she maintain ? 6
- (b) Find the angle between the vectors  $x = [8, -20, 4]$  and  $y = [6, -15, 3]$ . Hence or otherwise comment whether  $x$  and  $y$  are parallel or not. 6
- (c) Using Newton's Second Law of Motion, find the resultant sum of forces on a 30 kg object in a three-dimensional coordinate system undergoing an acceleration of  $6 \text{ m/sec}^2$  in the direction of the vector  $[-2, 3, 1]$ . 6

P.T.O.

2. (a) If A and B are row equivalent matrices, then what is the relationship between Row Space of A and Row Space of B ? Also, determine whether the vector  $[2, 2, -3]$  is in the row space of the matrix : 6.5

$$A = \begin{bmatrix} 4 & -1 & 2 \\ -2 & 3 & 5 \\ 6 & 1 & 9 \end{bmatrix}$$

- (b) Find all eigenvalues corresponding to the given matrix. Also, express each eigenspace as a set of linear combinations of fundamental eigenvectors. The given matrix is : 6.5

$$B = \begin{bmatrix} 12 & -51 \\ 2 & -11 \end{bmatrix}$$

- (c) (i) Show that the set  $\phi$  of real-valued functions,  $f$ , defined on the interval  $[0, 1]$  such that  $f\left(\frac{1}{2}\right) = 1$ , is not a vector space under the usual operations of function addition and scalar multiplication.
- (ii) Define subspace of a vector space. Prove or disprove that the set of 2-vectors of the form  $[a, 2a]$  is a subspace of  $\mathbf{R}^2$  under the usual vector operations, where  $a$  represents an arbitrary real number. 2+4.5
3. (a) Use the Simplified Span Method to find a simplified general form for all the vectors in Span (S), where  $S = \{[1, 1, 0], [2, -3, -5]\}$  is a subset of  $\mathbf{R}^3$ . 6

(b) Find if the homogeneous system  $Ax = 0$ , where :

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 0 & 1 & 0 \\ 1 & 0 & 3 \end{bmatrix},$$

has a non-trivial solution, using rank of A.

6

(c) Show that the set  $B = \{[1, -2, 1], [5, -3, 0]\}$  is the maximal linearly independent subset of  $S = \{[1, -2, 1], [3, 1, -2], [5, -3, 0], [5, 4, -5], [0, 0, 0]\}$ . Calculate  $\dim(\text{Span}(S))$ . Is  $\text{Span}(S) = \mathbf{R}^3$  ? Why or why not ?

6

4. (a) State the Dimension Theorem. Verify it for the linear transformation  $L : \mathbf{R}^3 \rightarrow \mathbf{R}^2$  given by :

6.5

$$L \left( \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} \right) = \begin{bmatrix} 3 & 2 & 11 \\ 2 & 1 & 8 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}.$$

(b) Find the matrix for the linear transformation L with respect to the standard bases for  $P_3$  and  $\mathbf{R}^3$ , where  $L : P_3 \rightarrow \mathbf{R}^3$  is defined by :


$$L(a_3x^3 + a_2x^2 + a_1x + a_0) = [a_0 + a_1, 2a_2, a_3 - a_0].$$

Use this matrix to compute  $L(5x^3 - x^2 + 3x + 2)$  by matrix multiplication.

6.5

P.T.O.

- (c) For the adjoining graphic, use ordinary coordinates in  $\mathbf{R}^2$  to find the new vertices after performing a rotation about the point (12, 6) through an angle  $\theta = 90^\circ$ . Then sketch the figure that would result from this movement. 6.5



5. (a) Let  $V$  be  $P_1$  and let  $S = \{v_1, v_2\}$  and  $T = \{w_1, w_2\}$  be ordered bases for  $P_1$ , where  $v_1 = t$ ,  $v_2 = t - 3$ ,  $w_1 = t - 1$ ,  $w_2 = t + 1$ . Compute the transition matrix  $Q_{T \leftarrow S}$  from the  $S$ -basis to the  $T$ -basis. 6
- (b) Define a linear operator. Show that the mapping  $f: P_n \rightarrow P_n$  defined by :
- $$f(p) = p + p'$$
- is a linear operator on  $P_n$ . 6
- (c) Is the linear transformation  $L: \mathbf{R}^3 \rightarrow \mathbf{R}^3$  given by :
- $$L([x, y, z]) = [2x, x + y + z, -y]$$
- one-to-one? Also, check whether it is onto or not. 6
6. (a) Show that the vector space  $\mathbf{R}^4$  is isomorphic to the vector space  $P_3$ , using the transformation  $L: \mathbf{R}^4 \rightarrow P_3$  given by  $L([a, b, c, d]) = ax^3 + bx^2 + cx + d$ . 6.5
- (b) Verify that the set  $B = \{[1, 0, -1], [-1, 4, -1], [2, 1, 2]\}$  is an orthogonal basis for  $\mathbf{R}^3$ . Obtain from  $B$  an orthonormal basis for  $\mathbf{R}^3$ . 6.5
- (c) Find the orthogonal complement  $W^\perp$  of the subspace  $W = \{[a, b, 0] | a, b \in \mathbf{R}\}$  of  $\mathbf{R}^3$ . Verify that  $\dim(W) + \dim(W^\perp) = \dim(\mathbf{R}^3)$ . 6.5