[This question paper contains 4 printed pages.]

Sr. No. of Question Paper: 106 E Your Roll No.....

Unique Paper Code : 236351

Name of the Course : B.A. Programme - Operational Research

Name of the Paper : Mathematical Programming

Semester : III

Duration: 3 Hours Maximum Marks: 75

Instructions for Candidates

- 1. Write your Roll No. on the top immediately on receipt of this question paper.
- 2. Attempt any five questions.
- 3. Use of non-programmable calculator is allowed.
- 1. (a) What is an unconstrained optimization problem? Define the concept of stationary point and relative extrema. (7)
 - (b) Determine the relative maximum and minimum (if any) of the following function:

$$f(x_1, x_2, x_3) = x_1 + 2x_2 + x_2x_3 - x_1^2 - x_2^2 - x_3^2$$
 (8)

- 2. (a) Define a convex function and prove that a positive linear combination of convex functions is again a convex function. (6)
 - (b) Use the Kuhn-Tucker conditions to solve the following non-linear programming problem:

Maximize
$$Z = 2x_1 - x_1^2 + x_2$$

Subject to constraints

$$2x_{1} + 3x_{2} \le 6$$

$$2x_{1} + x_{2} \le 4$$

$$x_{1} \ge 0, x_{2} \ge 0$$
(9)

3. Use the Wolfe's method for solving the following quadratic problem

Maximize
$$Z = 2x_1 + 3x_2 - 2x_1^2$$

Subject to $x_1 + 4x_2 \le 4$
 $x_1 + x_2 \le 2$
 $x_1, x_2 \ge 0$ (15)

- 4. (a) What is an Integer Programming Problem? Give some applications of integer programming problems. (7)
 - (b) Solve the following non-linear programming problem using the method of Lagrangian multipliers.

Maximize
$$Z = 4x_1 + 6x_2 - 2x_1^2 - 2x_1x_2 - 2x_2^2$$

subject to $x_1 + 2x_2 = 2$, and $x_1, x_2 \ge 0$. (8)

5. Use the branch and bound technique to solve the following Integer programming problem

Maximize
$$Z = x_1 + x_2$$

Subject to $2x_1 + 5x_2 \le 16$

$$6x_1 + 5x_2 \le 30$$

 $x_1, x_2 \ge 0$, and are integers.

Given non-integer optimum solution is: $x_1 = 7/2$, $x_2 = 9/5$ and Maximum Z = 53/10. (15)

6. (a) The following table provides a non-integer Optimum solution to IPP:

Maximize $Z = 4x_1 + 2x_2$

Subject to

$$x_1 + x_2 \le 7$$

$$2x_1 \le 11$$

$$2x_2 \leq 7$$

 $x_1, x_2 \ge 0$, and are integers.

	Optimum feasible non-integer solution						
ļ	c _j	4	2	0	0	0	
Св	X _B	x _i	x ₂	sı	s ₂	S ₃	ь
1	x,	1	0	1	0	-0.5	3.5
0	s ₂	0	0	-2	1	1	4
2	x ₂	0	1	0	0	0.5	3.5

Find the integer solution by using Gomory's fractional cut algorithm for mixed integer programming problem. (8)

(b) Prove that the following function is convex:

$$f(x_1, x_2) = 2x_1^2 + 3x_1x_2 + 2x_2^2 - 10x_1 - 10x_2, (x_1, x_2) \in \mathbb{R}^2.$$
 (7)

- 7. Write a short note on :-
 - (a) Branch and Bound Algorithm
 - (b) Bordered Hessian matrix
 - (c) Convex and Concave functions (15)